



Pitta Dosha and Human Thermoregulation: A Critical Synthesis of Ayurvedic Physiology and Modern Evidence Narrative Review

Apeksha Panwar^{1*}

¹Associate Professor Department of Kriya Sharir Himalayiya Ayurvedic Medical College and Hospital, Dehradun

* Corresponding author. E-mail: apekshanegi92@gmail.com

<https://doi.org/10.48165/irjay.2026.9.05.07>

ARTICLE INFO

Article history

Received: 22.07.2026

Revised: 11.08.2026

Accepted: 09.09.2026

Keywords

Pitta

Thermoregulation

Kriya Sharir

Prakriti

Body temperature

Thermal imaging

Agni

Sweat

Bhrajaka

Ayurveda

ABSTRACT

In Ayurvedic physiology, *Pitta* stands as the principal force governing transformation, metabolic activity and the generation of bodily heat. Classical descriptions repeatedly link its balanced state with stable internal temperature and its disturbance with intolerance to heat, excessive perspiration and a subjective sense of burning. Contemporary physiology, by contrast, accounts for temperature regulation through hypothalamic control, adjustments in metabolic output, changes in skin blood flow and evaporative cooling. Although the two systems appear to describe overlapping phenomena, few attempts have been made to place classical *Pitta* attributes side by side with measurable thermoregulatory variables. The present narrative review gathers material from the major *Samhitas* and their commentaries and from modern databases in order to examine this interface. Classical statements concerning *Ushna*, *Tikshna* and *Bhrajaka Pitta* are set against data on core and skin temperature, resting metabolic rate, sweat characteristics, infrared imaging and responses to heat stress. Available observations, though still limited in scale, point toward higher baseline temperatures, more abundant sweating and lower heat tolerance among individuals of *Pitta*-dominant constitution. Continuous temperature recording and thermal imaging offer practical means of testing these associations further. Important gaps remain, especially the absence of adequately controlled experimental work. Closer dialogue between the two traditions can strengthen the empirical base of *Kriya Sharir* and open useful avenues for individualised approaches to thermal comfort and heat-related risk.

INTRODUCTION

Stable core temperature is a prerequisite for the orderly function of enzymes, membrane transport and neural signalling. Modern accounts place the primary regulatory centre in the preoptic region of the hypothalamus, where inputs from deep and superficial thermoreceptors are integrated and matched against a defended set-point. Effector responses then adjust heat production through metabolic and muscular means, redistribute blood flow at the skin surface, and, when necessary, initiate sweating so that evaporative loss can offset the

heat load.¹²

Ayurveda approaches the same biological requirement through the doctrine of the three *Doshas*. *Pitta* is singled out as the agency of heat and transformation. Its cardinal qualities—hot, sharp, liquid and mobile—are repeatedly invoked to explain digestion, tissue metabolism, coloration of the blood and skin, and the maintenance of bodily warmth.^{1–3} When *Pitta* remains within normal limits, temperature is said to stay comfortable and digestion proceeds smoothly. When it increases, the individual experiences internal heat, burning sensations, heavy sweating and marked discomfort in warm surroundings. When it declines, coldness and sluggish metabolism appear.

These traditional observations raise a natural question: how far do they correspond to variables that can be measured with present-day instruments? A number of conceptual papers and a handful of small observational studies have noted differences in temperature and sweating according to *Prakriti*, yet a sustained, critical examination of the entire interface has not been published.^{4–6} The review that follows therefore sets out classical statements about *Pitta* in relation to heat, places them beside modern physiological parameters, surveys the limited empirical material that exists, considers possible underlying mechanisms, and outlines the clinical and research implications that follow for the field of *Kriya Sharir*.

CLASSICAL BACKGROUND: PITTA AND THE REGULATION OF HEAT

Linguistic and Conceptual Roots

The word *Pitta* is connected with the verbal root that denotes heating or burning. Early medical writers treat it as the bodily expression of the transformative power otherwise known as *Agni*. Charaka and Sushruta list its qualities as hot, sharp, fluid, spreading, light and slightly odorous. Of these, the quality of heat is given primacy because it sustains the metabolic fire that keeps the organism alive. Commentators repeatedly stress that the extinction of this internal heat is incompatible with continued life.^{1,2}

The relationship between *Pitta* and *Agni* is presented as reciprocal. When *Pitta* is well balanced, digestive and tissue fires operate efficiently and temperature remains steady. Excess *Pitta* intensifies heat production and produces the familiar signs of burning and intolerance; deficiency reduces both metabolic drive and body warmth. This classical pairing supplies a ready conceptual link with contemporary measurements of resting energy expenditure and thermogenesis.^{1–3,5}

The Five Functional Divisions of Pitta

Traditional physiology divides *Pitta* into five subtypes, each assigned a seat and a distinctive role. *Pachaka Pitta*, located in the stomach and upper intestine, is responsible for the primary digestion of food and is regarded as the chief source of metabolic heat arising from the gut. *Ranjaka Pitta*, associated with the liver and blood, contributes to the colouring of *Rakta* and to the distribution of heat throughout the circulation. *Sadhaka Pitta*, linked with the heart and mental faculties, influences emotional tone; anger and intense mental effort can themselves raise the sensation of heat.

Alochaka Pitta governs vision and is only indirectly related to systemic temperature. *Bhrajaka Pitta*, residing in the skin, is of special interest for thermoregulation. Classical authorities credit it with the maintenance of normal surface temperature, the glow of the complexion, and the processing of substances applied to the skin. In effect, *Bhrajaka Pitta* represents the traditional understanding of the skin as the principal organ of heat exchange with the external environment.^{1–3}

Clinical Markers of Pitta Balance and Imbalance

In a state of equilibrium, body warmth is moderate, sweating matches the demands of climate and activity, and the skin remains clear. When *Pitta* increases, the subject reports a feeling of internal heat, burning in the eyes, palms or soles, free sweating that may carry a noticeable odour, marked thirst, and clear intolerance of hot weather or pungent food. The skin may appear flushed and the hair may grey or thin earlier than expected. When *Pitta* diminishes, coldness of the body, weak digestion and loss of lustre become prominent. Sweat itself is described as a waste product of fat tissue that stands in an abode-content relationship with *Pitta*; its formation and release serve both excretory and cooling functions. Classical writers already noted that the amount and character of sweat differ according to constitution—an observation that invites direct comparison with modern sweat-rate and composition studies.^{1–4}

BRIEF OUTLINE OF CONTEMPORARY THERMOREGULATORY PHYSIOLOGY

Present-day understanding places the main controller in the preoptic hypothalamus. Afferent signals from core and skin thermoreceptors are compared with a regulated reference value. Any deviation activates appropriate effectors. Heat production rises through increases in basal metabolism, the energetic cost of food processing, muscular activity and, when required, shivering or the recruitment of brown adipose tissue. Heat is conserved by constriction of skin vessels or dissipated by their dilation. Once ambient temperature or internal heat load exceeds the capacity of dry heat loss, eccrine sweating provides the principal route of evaporative cooling.^{12–16}

A range of techniques now permits quantitative assessment of these processes. Core temperature can be tracked continuously by specialised sensors; skin temperature can be mapped at multiple sites or imaged by infrared cameras; metabolic rate can be estimated by

Table 1: Provisional mapping of Pitta characteristics onto thermoregulatory measures

Classical descriptor	Corresponding modern variable	Strength of present evidence
<i>Ushna</i> (heat quality)	Core and mean skin temperature; resting metabolic rate	Limited observational data
<i>Sweda</i> (perspiration)	Sweat rate and compositional profile	Classical statements plus sparse measurements
<i>Daha</i> and heat intolerance	Thermal comfort scores; heat-strain indices	Consistent clinical reports
<i>Tikshna</i> (sharp quality)	Speed of thermal response	Very few data
<i>Bhrajaka Pitta</i>	Cutaneous blood flow; infrared temperature patterns	Early clinical application
Link with Agni	Resting energy expenditure; substrate use	Indirect support from Prakriti studies

indirect calorimetry; sweat can be collected and analysed for volume and solute content; and standardised heat-exposure protocols allow measurement of tolerance thresholds and cardiovascular strain. These methods supply the concrete endpoints against which classical qualitative statements can be examined.^{13,17,18}

CORRESPONDENCE BETWEEN CLASSICAL ATTRIBUTES AND MEASURABLE VARIABLES

The following table summarises a working correspondence between traditional descriptors and modern parameters. The alignment is offered as a guide for further inquiry rather than as a claim of exact equivalence.

The quality of heat most readily invites comparison with measured temperature and metabolic rate. Descriptions of sweating correspond to volume and character of sweat. Subjective burning and intolerance align with comfort ratings and objective strain under heat load. The cutaneous seat of *Bhrajaka Pitta* suggests a natural focus on skin blood flow and thermal imaging. The classical association of *Pitta* with digestive and tissue fire points toward energy-expenditure measurements. Each of these links remains provisional and requires systematic testing.

SURVEY OF AVAILABLE OBSERVATIONAL MATERIAL

Direct experimental comparisons of continuous thermoregulatory records across well-characterised *Prakriti* groups are still scarce. What exists is largely observational and modest in scale, yet the pattern that emerges is consistent enough to warrant attention.

Studies that have recorded ordinary vital signs in relation to constitution report modestly higher average body temperatures among subjects classified as *Pitta*-predominant or as mixed types containing a strong *Pitta* component. One investigation conducted in healthy infants detected statistically detectable differences across *Prakriti* categories, with higher readings more often associated with *Pitta* influence.^{6,19} Although sample sizes have been small and assessment methods not fully standardised, the direction of the findings accords with classical expectation.

Descriptions of sweating likewise show continuity between traditional and modern accounts. *Pitta*-dominant individuals are regularly said to sweat more freely and to produce sweat that is warmer and sometimes more noticeable in odour. Sparse contemporary measurements of sweat rate, together with genetic indications of more rapid drug metabolism in *Pitta* types, are compatible with a higher baseline metabolic output.^{4,7,8} The traditional remark concerning the odour of sweat may ultimately relate to differences in apocrine activity or in the bacterial processing of sweat constituents—possibilities that have not yet been examined systematically.

Infrared thermography has begun to find a place in some Ayurvedic clinics as a means of visualising regional heat distribution. Controlled comparisons that stratify participants by rigorously determined *Prakriti* are lacking, yet the method is non-invasive and capable of generating quantitative maps of surface temperature.^{17,18} Wearable sensors that log temperature continuously further enlarge the scope for ambulatory study. Heat intolerance itself remains a recurrent clinical observation among *Pitta*-predominant persons and matches modern notions of reduced tolerance and greater physiological cost under thermal stress. Formal heat-challenge trials stratified by constitution have not, however, been reported.

HYPOTHETICAL BIOLOGICAL PATHWAYS

Several physiological routes could explain the associations sketched above. Elevated sympathetic tone and higher circulating catecholamines would be expected to raise both metabolic rate and skin blood flow. Enhanced activity of mitochondrial uncoupling proteins or of the thyroid axis could increase non-shivering heat production. Low-grade inflammatory or oxidative processes, themselves influenced by diet and lifestyle factors that also affect *Pitta*, might lower the threshold for the subjective sensation of heat. At the skin surface, variation in vascular reactivity or in the density and responsiveness of sweat glands could account for differences traditionally ascribed to *Bhrajaka Pitta*.^{7,10,11,16}

None of these suggestions has yet been tested under controlled conditions. Progress will require combined approaches that include continuous autonomic recording, targeted molecular assays, and thermal challenges performed in participants whose constitution has been assessed with reproducible instruments.

IMPLICATIONS FOR PRACTICE AND TRANSLATION

If constitution-related differences in thermal physiology can be confirmed, several practical consequences follow. Advice on clothing, indoor climate, timing of outdoor work and exercise intensity could be adjusted according to *Dosha* predominance. Individuals with a strong *Pitta* component might be counselled to adopt cooling measures earlier in the season or during periods of high physical demand. Continuous temperature monitors and thermal imaging could serve as objective markers for tracking response to traditional cooling regimens—dietary adjustments, specific herbal preparations, oil massage with cooling media, and selected purification procedures. In occupational and sporting contexts, early identification of persons at greater risk of heat illness could guide preventive strategies. At the same time, a clearer physiological understanding of classical cooling therapies may suggest new adjunctive options for conditions marked by excessive heat generation or impaired heat loss.^{20–23}

OUTSTANDING QUESTIONS AND RESEARCH PRIORITIES

Several concrete deficiencies limit present confidence in the proposed correlations:

1. Large, multi-centre investigations that pair validated *Prakriti* instruments with continuous core and multi-site skin temperature records under both resting and heat-loaded conditions are still missing.
2. Standardised heat-exposure protocols that classify participants by *Dosha* and report objective indices of strain, sweating threshold and recovery have not been published.
3. Infrared and wearable temperature data have yet to be validated against classical signs of *Pitta* and against clinical response to cooling measures.
4. Molecular, autonomic and endocrine profiles linked to *Pitta* status remain largely unexplored.
5. Adequately powered trials that test classical *Pitta*-pacifying interventions with pre-specified thermoregulatory endpoints are scarce.

Closing these gaps will demand sustained cooperation among Ayurvedic scholars, physiologists, engineers and clinical investigators. Standardised assessment tools, transparent protocols and balanced reporting of results will be essential if the field is to move from plausible correspondence to reliable evidence.

CONCLUDING REMARKS

Classical statements about *Pitta* supply a coherent qualitative account of how individuals differ in the production, distribution and loss of heat. Modern physiological techniques offer the means to test and refine that account. The limited observations so far available are broadly consistent with traditional expectations, especially regarding baseline temperature, sweating and heat intolerance. At the same time, the evidence remains thin and largely exploratory. A two-way exchange—letting classical insight shape precise questions and letting quantitative measurement correct and extend traditional concepts—can deepen the scientific grounding of *Kriya Sharir* and contribute practical tools for personalised thermal care. Systematic work along these lines is both feasible and needed.

ACKNOWLEDGEMENTS

The author thanks the classical physicians whose observations form the starting point of this discussion and the contemporary researchers whose measurements make critical comparison possible.

AUTHOR CONTRIBUTIONS

The author took part in framing the review, locating and assessing sources, interpreting the material and preparing the final text.

FUNDING

No external funding was received for this work.

ETHICAL STATEMENT

The article is a literature-based narrative review and involves neither human participants nor experimental animals; formal ethical approval is therefore not required.

CONFLICT OF INTEREST

The author declares no competing interests.

REFERENCES

- Sharma, P. V. (Ed.). (2014). *Charaka Samhita of Agnivesha*. Chaukhambha Orientalia.
- Sharma, P. V. (Ed.). (2013). *Sushruta Samhita of Sushruta*. Chaukhambha Visvabharati.
- Srikantha Murthy, K. R. (Trans.). (2012). *Ashtanga Hridaya of Vagbhata*. Chaukhambha Krishnadas Academy.
- Verma, V., Gehlot, S., & Agrawal, S. (2019). Ayurveda insights on physiology of sweating and thermoregulation. *Journal of Natural Remedies*, 19(3), 111–120.
- Hankey, A. (2010). Establishing the scientific validity of Tri-dosha. *Ancient Science of Life*, 29(3), 6–18.
- Rotti, H., Raval, R., Anchan, S., et al. (2014). Determinants of Prakriti and its correlation with contemporary science. *Journal of Ayurveda and Integrative Medicine*, 5(3), 167–175.
- Prasher, B., Negi, S., Aggarwal, S., et al. (2008). Whole genome expression and biochemical correlates of extreme constitutional types. *Journal of Translational Medicine*, 6, 48.
- Ghodke, Y., Joshi, K., & Patwardhan, B. (2011). Ayurveda Prakriti type and CYP2C19 gene polymorphism. *Evidence-Based Complementary and Alternative Medicine*, 2011, 249528.
- Bhalerao, S., Deshpande, T., & Thatte, U. (2012). Prakriti and variations in platelet aggregation. *BMC Complementary and Alternative Medicine*, 12, 248.
- Dey, S., & Pahwa, P. (2014). Prakriti and its associations with metabolism, genetics and epigenetics. *Journal of Ayurveda and Integrative Medicine*, 5(1), 15–24.
- Govindaraj, P., Nizamuddin, S., Sharath, A., et al. (2015). Genome-wide analysis correlates Ayurveda Prakriti. *Scientific Reports*, 5, 15786.
- Romanovsky, A. A. (2018). The thermoregulation system and how it works. *Handbook of Clinical Neurology*, 156, 3–43.
- Kenny, G. P., & Jay, O. (2013). Thermometry, calorimetry and mean body temperature during heat stress. *Comprehensive Physiology*, 3(4), 1689–1719.
- Cramer, M. N., & Jay, O. (2016). Biophysical aspects of human thermoregulation during heat stress. *Autonomic Neuroscience*, 196, 3–14.
- Shibasaki, M., & Crandall, C. G. (2010). Mechanisms and controllers of eccrine sweating in humans. *Frontiers in Bioscience*, 2, 685–696.
- Charkoudian, N. (2003). Skin blood flow in adult human thermoregulation. *Mayo Clinic Proceedings*, 78(5), 603–612.
- Fernandez-Cuevas, I., Marins, J. C., Lastras, J. A., et al. (2015). Classification of factors influencing infrared thermography in humans. *Infrared Physics & Technology*, 71, 28–55.
- Ring, E. F. J., & Ammer, K. (2012). Infrared thermal imaging in medicine. *Physiological Measurement*, 33(3), R33–R46.
- Tiwari, S., Gehlot, S., & Tiwari, S. K. (2011). Effect of temperature and seasonal variation on Prakriti. *AYU*, 32(2), 211–215.
- Patwardhan, B., Mutalik, G., & Tillu, G. (2015). *Integrative approaches for health*. Academic Press.
- Hankey, A. (2005). The scientific value of Ayurveda. *Journal of Alternative and Complementary Medicine*, 11(2), 221–225.
- Rastogi, S. (2010). Building bridges between Ayurveda and modern science. *International Journal of Ayurveda Research*, 1(1), 41–46.
- Joshi, K., Ghodke, Y., & Shintre, P. (2010). Traditional medicine and genomics. *Journal of Ayurveda and Integrative Medicine*, 1(1), 26–32.